






## NRBP2 Antibody

<b>Product Code</b>	CSB-PA878883LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9NSY0
Immunogen	Recombinant Human Nuclear receptor-binding protein 2 protein (12-258AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, IHC, IF, IP; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200, IP:1:200-1:2000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Nuclear receptor-binding protein 2 (Transformation-related gene 16 protein) (TRG-16), NRBP2
Immunogen Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Target Names	NRBP2
Image	Immunohistochemistry of paraffin-embedded



Immunohistochemistry of paraffin-embedded human endometrial cancer using CSB-PA878883LA01HU at dilution of 1:100
















Immunohistochemistry of paraffin-embedded human adrenal gland tissue using CSB-PA878883LA01HU at dilution of 1:100



Immunofluorescence staining of Hela cells with CSB-PA878883LA01HU at 1:133, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunoprecipitating NRBP2 in A549 whole cell lysate

Lane 1: Rabbit control IgG (1µg) instead of CSB-PA87883LA01HU in A549 whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)

Lane 2: CSB-PA878883LA01HU (6µg) + A549 whole cell lysate (500µg)

Lane 3: A549 whole cell lysate (10µg)