



# PFDN1 Antibody Pair

|                            |   |
|----------------------------|---|
| <b>Product Code</b>        | CSB-EAP01485  |
| <b>Species Reactivity</b>  | Human,Rat,Mouse,Guinea pig,Bovine,Pig,Dog,Goat  |
| <b>Tested Applications</b> | S-ELISA   |
| <b>Form</b>                | Liquid  |
| <b>Alias</b>               | PFDN1,PFD1  |
| <b>Product Type</b>        | Antibody Pairs  |
| <b>Immunogen Species</b>   | Homo sapiens (Human)  |
| <b>Gene Names</b>          | PFDN1   |
| <b>Protein Names</b>       | Prefoldin subunit 1   |
| <b>Notes</b>               | We recommend using the capture antibody at a concentration of 0.5ug/ml and the detection antibody at a concentration of 0.5ug/ml.Optimal dilutions should be determined experimentally by the researcher. |
| <b>Host</b>                | Capture: Rabbit<br>Detection: Rabbit  |
| <b>Components</b>          | Capture: CSB-EAP01485C<br>Detection: CSB-EAP01485D(Biotin)<br>Reagents are sufficient for at least 5 x 96 well plates using recommended protocol.   |
| <b>Storage-Buffer</b>      | Capture: 50% Glycerol, 0.01M PBS, PH 7.4<br>Detection: 50% Glycerol, 0.01M PBS, PH 7.4  |