



Recombinant Mouse Dipeptidyl peptidase 4 (Dpp4), partial


Product Code	CSB-CF007139MO
Relevance	<p>Cell surface glycoprotein receptor involved in the costimulatory signal essential for T-cell receptor (TCR)-mediated T-cell activation. Acts as a positive regulator of T-cell coactivation, by binding at least ADA, CAV1, IGF2R, and PTPRC. Its binding to CAV1 and CARD11 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner. Its interaction with ADA also regulates lymphocyte-epithelial cell adhesion. In association with FAP is involved in the pericellular proteolysis of the extracellular matrix (ECM), the migration and invasion of endothelial cells into the ECM. May be involved in the promotion of lymphatic endothelial cells adhesion, migration and tube formation. When overexpressed, enhanced cell proliferation, a process inhibited by GPC3. Acts also as a serine exopeptidase with a dipeptidyl peptidase activity that regulates various physiological processes by cleaving peptides in the circulation, including many chemokines, mitogenic growth factors, neuropeptides and peptide hormones. Removes N-terminal dipeptides sequentially from polypeptides having unsubstituted N-termini provided that the penultimate residue is proline.</p>
Abbreviation	Recombinant Mouse Dpp4 protein, partial
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P28843
Product Type	Transmembrane Protein
Immunogen Species	Mus musculus (Mouse)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	<p>SKDEAAADSRRTYSLADYLKSTFRVKSYSYLWWVSDFEYLYKQENNILLNNAEH GNSSIFLENSTFESFGYHSVSPDRLFVLLLEYNYVKQWRHSYASYNIIYDVNKR QLITEEKIPNNTQWITWSPEGHKLAYVWKNDIYVKVEPHLPSHRITSTGEENVIIY NGITDWVYEEEVFGAYSALWWSPNNTFLAYAQFNDTGVPLEIYSFYSDLSLQY PKTVWIPYPKAGAVNPTVKFFIVNIDSLSSSSSAAPIQIPAPASVARGDHLYLCDV VWATEERISLQWLRRIQNYSVMAICDYDKINLTWNCNPSEQQHVMSTTGWVG RFRPAEPHFTSDGSSFYKIIISDKDGYKHICHFPKDKKDCFTITKGAWEVISIEAL TSDYLYYISNQYKEMPGGRNLYKIQLTDHTNVKCLSCDLNPERCQYYAVSFSK EAKYYQLGCWGPGLPLYTLHRSTDHKLRLVLEDNSALDRMLQDVQMPSKCLD FIVLNETRFWYQMILPPHFDKSKKYPLLLDVYAGPCSQKADASFRLNWATYLA STENIIVASFDGRGSGYQGDKIMHAINRRRLGTLEVEDQIEAARQFVKMGFVDSK RVAIWGWSYGGYVTSMLVLSGSGVFKCGIAPVSRWEYYSVYTERYMGL PIPEDNLDHYRNSTVMSRAEHFKQVEYLLIHGTADDNVHFQQSAQISKALVDA</p>



GVDFQAMWYTDEDHGIASSTAHQHIYSHMSHFLQQCFSLH

Research Area	Immunology
Source	in vitro E.coli expression system
Target Names	Dpp4
Protein Names	Dipeptidyl peptidase IV Short name:DPP IVT-cell activation antigen CD26Thymocyte-activating molecule Short name:THAMCD_antigen: CD26Cleaved into the following 2 chains:Dipeptidyl peptidase 4 membrane form Alternative name(s):Dipeptidyl peptidase I
Expression Region	29-760aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal 6xHis-SUMO-tagged
Mol. Weight	100.5kDa
Protein Length	Extracellular Domain

Image



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.