




# CENPK Antibody

<b>Product Code</b>	CSB-PA005214ESR1HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q9BS16
<b>Immunogen</b>	Recombinant Human Centromere protein K protein (1-269AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IHC, IP; Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200, IP:1:200-1:2000
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Purification Method</b>	Antigen Affinity Purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Centromere protein K (CENP-K) (Interphase centromere complex protein 37) (Protein AF-5alpha) (p33), CENPK, ICEN37
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Epigenetics and Nuclear Signaling
<b>Target Names</b>	CENPK

## Image



### Western blot

All lanes: CENPK antibody at 3.98μg/ml

Lane 1: PC-3 whole cell lysate


Lane 2: 293T whole cell lysate

Secondary


Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 32 kDa


Observed band size: 32 kDa



Immunohistochemistry of paraffin-embedded human skin tissue using CSB-PA005214ESR1HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human liver tissue using CSB-PA005214ESR1HU at dilution of 1:100



Immunoprecipitating CENPK in HeLa whole cell lysate  
 Lane 1: Rabbit control IgG instead of CSB-PA005214ESR1HU in HeLa whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)  
 Lane 2: CSB-PA005214ESR1HU (8μg) + HeLa whole cell lysate (500μg)  
 Lane 3: HeLa whole cell lysate (10μg)