




# FBXO22 Antibody

<b>Product Code</b>	CSB-PA854115ESR1HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q8NEZ5
<b>Immunogen</b>	Synthesized peptide derived from human FBX22 AA range: 182-232
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human, Mouse
<b>Tested Applications</b>	ELISA, WB, IHC; Recommended dilution: WB:1:1000-1:3000, IHC:1:100-1:300
<b>Form</b>	Liquid
<b>Storage Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Purification Method</b>	Antigen Affinity Purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Product Type</b>	Polyclonal Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Cell Biology
<b>Target Names</b>	FBXO22

## Image



### Western Blot

Positive WB detected in: Mouse Brain tissue lysate(20µg)

All lanes: FBXO22 antibody at 1:1000


Secondary

Goat polyclonal to rabbit IgG at 1/20000 dilution

Predicted band size: 45kDa

Observed band size: 45 kDa

Exposure time: 10s



### Western Blot

Positive WB detected in: HeLa whole cell lysate(20µg), MCF7 whole cell lysate(20µg), HepG2 whole cell lysate(20µg), 293 whole cell lysate(20µg), JK whole cell lysate(20µg), K562 whole cell lysate(20µg), A549 whole cell lysate(20µg), A431 whole cell lysate(20µg), U251 whole cell lysate(20µg)

All lanes: FBXO22 antibody at 1:1000


Secondary

Goat polyclonal to rabbit IgG at 1/20000 dilution


Predicted band size: 45 kDa



Observed band size: 45 kDa  
Exposure time: 10s



IHC image of CSB-PA854115ESR1HU diluted at 1:200 and staining in paraffin-embedded human liver tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB. Secondary antibody only control: uses 1% BSA instead of primary antibody



IHC image of CSB-PA854115ESR1HU diluted at 1:200 and staining in paraffin-embedded human tonsil tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB. Secondary antibody only control: uses 1% BSA instead of primary antibody

## Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.