




# EMCN Antibody

<b>Product Code</b>	CSB-PA007632ESR1HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q9ULC0
<b>Immunogen</b>	Recombinant Human Endomucin protein (19-135AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IHC; Recommended dilution: WB:1:1000-1:3000, IHC:1:50-1:200
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Purification Method</b>	Antigen Affinity Purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Product Type</b>	Polyclonal Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Signal Transduction
<b>Target Names</b>	EMCN

## Image



### Western Blot

Positive WB detected in: HEK293 whole cell lysate(20µg), 293T whole cell lysate(20µg), PC-3 whole cell lysate(20µg)

All lanes: FFAR3 antibody at 1:1000


Secondary

Goat polyclonal to rabbit IgG at 1/40000 dilution

Predicted band size: 27, 19 kDa

Observed band size: 37 kDa

Exposure time: 5s



IHC image of CSB-PA007632ESR1HU diluted at 1:50 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and



visualized using 0.05% DAB. Secondary antibody only control: uses 1% BSA instead of primary antibody

---

**Usage**

For Research Use Only. Not for use in diagnostic or therapeutic procedures.