




EIF3K Antibody

Product Code	CSB-PA866204ESR1HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9UBQ5
Immunogen	Recombinant Human Eukaryotic translation initiation factor 3 subunit K protein (1-218AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC, IP; Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200, IP:1:200-1:2000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Purification Method	Antigen Affinity Purified
Isotype	IgG
Clonality	Polyclonal
Alias	Eukaryotic translation initiation factor 3 subunit K (eIF3k) (Eukaryotic translation initiation factor 3 subunit 12) (Muscle-specific gene M9 protein) (PLAC-24) (eIF-3 p25) (eIF-3 p28), EIF3K, EIF3S12
Immunogen Species	Homo sapiens (Human)
Research Area	Neuroscience
Target Names	EIF3K

Image



Western blot

All lanes: EIF3K antibody at 1.72μg/ml

Lane 1: THP-1 whole cell lysate

Lane 2: HT29 whole cell lysate

Lane 3: 293T whole cell lysate


Lane 4: Hela whole cell lysate

Secondary


Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 26, 25 kDa


Observed band size: 26 kDa



Immunohistochemistry of paraffin-embedded human liver cancer using CSB-PA866204ESR1HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human kidney tissue using CSB-PA866204ESR1HU at dilution of 1:100



Immunoprecipitating EIF3K in HeLa whole cell lysate

Lane 1: Rabbit control IgG instead of (1μg) instead of CSB-PA866204ESR1HU in HeLa whole cell lysate. For western blotting, a HRP-conjugated anti-rabbit IgG, specific to the non-reduced form of IgG was used as the Secondary antibody (1/50000)

Lane 2: CSB-PA866204ESR1HU (4μg) + HeLa whole cell lysate (500μg)

Lane 3: HeLa whole cell lysate (20μg)