



Recombinant Human Proteasome subunit beta type-10 (PSMB10)

Product Code	CSB-EP018877HU-B
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	P40306
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	>85% (SDS-PAGE)
Sequence	T TIAGLVFQDG VILGADTRAT NDSVVADKSC EKIHFIAPKI YCCGAGVAAD AEMTTRMVAS KMELHALSTG REPRVATVTR ILRQTLFRYQ GHVGASLIVG GVDLTGPQLY GVHPHGSYSR LPFTALGSGQ DAALAVLEDR FQPNTMLEAA QGLLVEAVTA GILGDLGSGG NVDACVITKT GAKLLRTLSS PTEPVKRSR YHFVPGTTAV LTQTVKPLTL ELVEETVQAM EVE
Source	E.coli
Target Names	PSMB10
Protein Names	Recommended name: Proteasome subunit beta type-10 EC= 3.4.25.1 Alternative name(s): Low molecular mass protein 10 Macropain subunit MECI-1 Multicatalytic endopeptidase complex subunit MECI-1 Proteasome MECI-1 Proteasome subu
Expression Region	40-273
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full Length of Mature Protein
Target Details	The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. Proteolytic processing is required to generate a mature subunit. Expression of this gene is induced by gamma interferon, and this gene product replaces catalytic subunit 2 (proteasome beta 7 subunit) in the immunoproteasome.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final



concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.