



# Recombinant Human Interleukin-15 (IL15)

<b>Product Code</b>	CSB-EP011593HU-B
<b>Storage</b>	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
<b>Uniprot No.</b>	P40933
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Purity</b>	>85% (SDS-PAGE)
<b>Sequence</b>	NWVNVISDLKKIEDLIQSMHIDATLYTESDVHPSCCKVTAMKCFLELQVISLESG DASIHDTVENLILANNLSLSSNGNVTESGCKECEEELEEKNIKEFLQSFVHIVQMFI NTS
<b>Research Area</b>	Metabolism
<b>Source</b>	E.coli
<b>Target Names</b>	IL15
<b>Protein Names</b>	Recommended name: Interleukin-15 Short name= IL-15
<b>Expression Region</b>	49-162aa
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	Full Length of Mature Protein
<b>Target Details</b>	This protein is a cytokine that regulates T and natural killer cell activation and proliferation. This cytokine and interleukine 2 share many biological activities. They are found to bind common hematopoietin receptor subunits, and may compete for the same receptor, and thus negatively regulate each other's activity. The number of CD8+ memory cells is shown to be controlled by a balance between this cytokine and IL2. This cytokine induces the activation of JAK kinases, as well as the phosphorylation and activation of transcription activators STAT3, STAT5, and STAT6. Studies of the mouse counterpart suggested that this cytokine may increase the expression of apoptosis inhibitor BCL2L1/BCL-x(L), possibly through the transcription activation activity of STAT6, and thus prevent apoptosis. Two alternatively spliced transcript variants of this gene encoding the same protein have been reported.
<b>Reconstitution</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
<b>Shelf Life</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life



of lyophilized form is 12 months at -20°C/-80°C.