








## **ARHGDIG** Antibody

<b>Product Code</b>	CSB-PA858727LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q99819
Immunogen	Recombinant Human Rho GDP-dissociation inhibitor 3 protein (1-225AA)
Raised In	Rabbit
Species Reactivity	Human, Mouse
Tested Applications	ELISA, WB, IHC, IF; Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200, IF:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Rho GDP-dissociation inhibitor 3 (Rho GDI 3) (Rho-GDI gamma), ARHGDIG
Immunogen Species	Homo sapiens (Human)
Research Area	Signal Transduction
Target Names	ARHGDIG
Imago	

**Image** 



Western blot

All lanes: ARHGDIG antibody at 2 μg/ml

Lane 1: Mouse brain tissue Lane 2: Mouse lung tissue

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 25 kDa Observed band size: 25, 50 kDa

















Immunohistochemistry of paraffin-embedded human brain tissue using CSB-PA858727LA01HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human placenta tissue using CSB-PA858727LA01HU at dilution of 1:100



Immunofluorescent analysis of MCF-7 cells using CSB-PA858727LA01HU at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Western Blot

Positive WB detected in: Hela whole cell lysate, 293 whole cell lysate, MCF-7 whole cell lysate, SH-SY5Y whole cell lysate, K562 whole cell lysate, A549 whole cell lysate

All lanes: ARHGDIG antibody at 1:2000

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 26 kDa Observed band size: 26 kDa