





## Recombinant Mouse Caspase-1 (Casp1), partial

| <b>Product Code</b> | CSB-BP004543MO  |
|---------------------|---|
| Storage             | Store at -20°C, for extended storage, conserve at -20°C or -80°C.   |
| Uniprot No.         | P29452  |
| Product Type        | Recombinant Protein   |
| Immunogen Species   | Mus musculus (Mouse)  |
| Purity              | >85% (SDS-PAGE)   |
| Source              | Baculovirus   |
| Target Names        | Casp1   |
| Protein Names       | Recommended name: Caspase-1 Short name= CASP-1 EC= 3.4.22.36 Alternative name(s): Interleukin-1 beta convertase Short name= IL-1BC Interleukin-1 beta-converting enzyme Short name= ICE Short name= IL-1 beta-co  |
| Expression Region   | -   |
| Notes               | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.   |
| Tag Info            | Tag type will be determined during the manufacturing process.   |
| Protein Length      | Partial   |
| Target Details      | This gene encodes a protein which is a member of the cysteine-aspartic acid   |
|                     | protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme. This gene was identified by its ability to proteolytically cleave and activate the inactive precursor of interleukin-1, a cytokine involved in the processes such as inflammation, septic shock, and wound healing. This gene has been shown to induce cell apoptosis and may function in various developmental stages. Studies of a similar gene in mouse suggest a role in the pathogenesis of Huntington disease. Alternative splicing of this gene results in five transcript variants encoding distinct isoforms. |
| Reconstitution      | protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme. This gene was identified by its ability to proteolytically cleave and activate the inactive precursor of interleukin-1, a cytokine involved in the processes such as inflammation, septic shock, and wound healing. This gene has been shown to induce cell apoptosis and may function in various developmental stages. Studies of a similar gene in mouse suggest a role in the pathogenesis of Huntington disease. Alternative splicing of this gene results in five transcript                                      |