



Recombinant Human NAD-dependent protein deacetylase sirtuin-6 (SIRT6)

Product Code	CSB-EP854057HU
Relevance	NAD-dependent protein deacetylase. Has deacetylase activity towards histone H3K9Ac and H3K56Ac. Modulates acetylation of histone H3 in telomeric chromatin during the S-phase of the cell cycle. Deacetylates histone H3K9Ac at NF-kappa-B target promoters and may down-regulate the expression of a subset of NF-kappa-B target genes. Acts as a corepressor of the transcription factor HIF1A to control the expression of multiple glycolytic genes to regulate glucose homeostasis. Required for genomic stability. Regulates the production of TNF protein. Has a role in the regulation of life span. Deacetylation of nucleosomes interferes with RELA binding to target DNA. May be required for the association of WRN with telomeres during S-phase and for normal telomere maintenance. Required for genomic stability. Required for normal IGF1 serum levels and normal glucose homeostasis. Modulates cellular senescence and apoptosis. On DNA damage, promotes DNA end resection via deacetylation of RBBP8. Has very weak deacetylase activity and can bind NAD ⁺ in the absence of acetylated substrate.
Abbreviation	Recombinant Human SIRT6 protein
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	Q8N6T7
Alias	Regulatory protein SIR2 homolog 6 SIR2-like protein 6
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	SVNYAAGLSPYADK GKCGLP EIFDPPEELERK V WELARLVWQSSSVVFHTGA GISTASGIPDFRGP HGVWTMEERGLAPKFD TTFESARPTQTHMALVQLERVGL LRFLVSQNV DGLHVRSGFPRDKLAELHGNMFVEECAKCKTQYVRD TVVGTM GLKATGRLCTVAKARGLRACRGELRDTILDWEDSLPDRDLALADEASRNADLS ITLGTSLQIRPSGNLPLATKRRGGRLVIVNLQPTKHDRHADLRIHGYVDEVMTR LMKHLGLEIPAWDGPVLERALPPLPRPPTPKLEPKESPTRINGSIPAGPKQE PCAQHNGSEPA SPKRERPTSPAPHRPPKRVKAKAVPS
Source	E.coli
Target Names	SIRT6
Protein Names	Recommended name: NAD-dependent protein deacetylase sirtuin-6 EC= 3.5.1.- Alternative name(s): Regulatory protein SIR2 homolog 6 SIR2-like protein 6
Expression Region	2-355aa




Notes Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Tag Info N-terminal GST-tagged

Mol. Weight 66.0kDa

Protein Length Full Length of Mature Protein

Image



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.