




# BTK Antibody

<b>Product Code</b>	CSB-PA002867LA01HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q06187
<b>Immunogen</b>	Recombinant Human Tyrosine-protein kinase BTK protein (254-443AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human, Mouse
<b>Tested Applications</b>	ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200, IF:1:50-1:200
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	>95%, Protein G purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Tyrosine-protein kinase BTK (EC 2.7.10.2) (Agammaglobulinemia tyrosine kinase) (ATK) (B-cell progenitor kinase) (BPK) (Bruton tyrosine kinase), BTK, AGMX1 ATK BPK
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Signal Transduction
<b>Target Names</b>	BTK

## Image



### Western Blot

Positive WB detected in: K562 whole cell lysate, HL60 whole cell lysate, Mouse heart tissue, Mouse brain tissue


All lanes: BTK antibody at 3.5µg/ml

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 77, 80 kDa

Observed band size: 77 kDa



Immunohistochemistry of paraffin-embedded human spleen tissue using CSB-PA002867LA01HU at dilution of 1:100



Immunofluorescent analysis of HeLa cells using CSB-PA002867LA01HU at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)