




# CD96 Antibody

<b>Product Code</b>	CSB-PA004971LA01HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	P40200
<b>Immunogen</b>	Recombinant Human T-cell surface protein tactile protein
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IF; Recommended dilution: WB:1:500-1:2000, IF:1:50-1:200
<b>Form</b>	Liquid
<b>Storage Buffer</b>	PBS with 0.01% thimerosal, 50% glycerol, pH7.3.
<b>Purification Method</b>	Antigen affinity purification
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Product Type</b>	Polyclonal Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Immunology
<b>Target Names</b>	CD96

## Image




### Western Blot


Positive WB detected in: MCF7 whole cell lysate(30µg), 293 whole cell lysate(30µg), HepG2 whole cell lysate(30µg), JK whole cell lysate(30µg), K562 whole cell lysate(30µg)  
All lanes: CD96 antibody at 1:1000

### Secondary

Goat polyclonal to rabbit IgG at 1/40000 dilution  
Predicted band size: 66,64 kDa  
Observed band size: 66,120 kDa  
Exposure time: 120s



Immunofluorescence staining of HepG2 cell with CSB-PA004971LA01HU at 1:110, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescence staining of HepG2 cell with 5% goat serum, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

## Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.