




# Recombinant Human X-ray repair cross-complementing protein 6 (XRCC6), partial

<b>Product Code</b>	CSB-RP016174h
<b>Relevance</b>	Single-stranded DNA-dependent ATP-dependent helicase. Has a role in chromosome translocation. The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner. It works in the 3'-5' direction. Binding to DNA may be mediated by XRCC6. Involved in DNA non-homologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination. The XRCC5/6 dimer acts as regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold. The XRCC5/6 dimer is probably involved in stabilizing broken DNA ends and bringing them together. The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step. Required for osteocalcin gene expression. Probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5' deoxyribose-5-phosphate at an abasic site near double-strand breaks. 5'-dRP lyase activity allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined. The XRCC5/6 dimer together with APEX1 acts as a negative regulator of transcription
<b>Abbreviation</b>	Recombinant Human XRCC6 protein, partial
<b>Storage</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
<b>Uniprot No.</b>	P12956
<b>Alias</b>	5'-deoxyribose-5-phosphate lyase Ku70 ;5'-dRP lyase Ku7070 kDa subunit of Ku antigen;ATP-dependent DNA helicase 2 subunit 1ATP-dependent DNA helicase II 70 kDa subunitCTC box-binding factor 75 kDa subunit ;CTC75 ;CTCBFDNA repair protein XR;CC6Lupus Ku autoantigen protein p70 ;Ku70Thyroid-lupus autoantigen ;TLAAX-ray repair complementing defective repair in Chinese hamster cells 6
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Purity</b>	≥ 90% as determined by SDS-PAGE.
<b>Sequence</b>	SYYKTEGDEEAEQQEENLEASGDYKYSGRDSLIFLVDASKAMFESQSEDELT PFDMSIQCIQSVYISKIISDRDLLAVVFYGTEDKDNSVNFKNIIYVLQELDNPGA KRILELDQFKGQQGQKRFQDMMGHGSDYSLSEVLWVCANLFSQVQFKMSHK RIMLFTNEDNPHGNDSAKASRARTKAGDLRDTGIFLDMHLKPKGGFDISLFYR DIIS
<b>Research Area</b>	Epigenetics and Nuclear Signaling



<b>Source</b>	E.coli
<b>Target Names</b>	XRCC6
<b>Expression Region</b>	6-222aa
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	N-terminal 6xHis-tagged
<b>Mol. Weight</b>	28.8kDa
<b>Protein Length</b>	Partial

**Image**


(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

<b>Reconstitution</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
<b>Shelf Life</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.