

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

## MET Antibody, HRP conjugated

Product CodeCSB-PA013714LB01HUStorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.P08581ImmunogenRecombinant Human Hepatocyte growth factor receptor protein (83-209AA)Raised InRabbitSpecies ReactivityHumanHumanELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHepatocyte growth factor receptor (HGF receptor) (EC 2.7.10.1) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), METTarget NamesMET		
Uniprot No.P08581Uniprot No.P08581ImmunogenRecombinant Human Hepatocyte growth factor receptor protein (83-209AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHepatocyte growth factor receptor (HGF receptor) (EC 2.7.10.1) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), METImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Product Code	CSB-PA013714LB01HU
ImmunogenRecombinant Human Hepatocyte growth factor receptor protein (83-209AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHepatocyte growth factor receptor (HGF receptor) (EC 2.7.10.1) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), METImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHepatocyte growth factor receptor (HGF receptor) (EC 2.7.10.1) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), METImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Uniprot No.	P08581
Species ReactivityHumanTested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHepatocyte growth factor receptor (HGF receptor) (EC 2.7.10.1) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), METImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Immunogen	Recombinant Human Hepatocyte growth factor receptor protein (83-209AA)
Tested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHepatocyte growth factor receptor (HGF receptor) (EC 2.7.10.1) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), METImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Raised In	Rabbit
FormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHepatocyte growth factor receptor (HGF receptor) (EC 2.7.10.1) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), METImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Species Reactivity	Human
ConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHepatocyte growth factor receptor (HGF receptor) (EC 2.7.10.1) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), METImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	<b>Tested Applications</b>	ELISA
Storage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHepatocyte growth factor receptor (HGF receptor) (EC 2.7.10.1) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), METImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Form	Liquid
Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHepatocyte growth factor receptor (HGF receptor) (EC 2.7.10.1) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), METImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Conjugate	HRP
IsotypeIgGClonalityPolyclonalAliasHepatocyte growth factor receptor (HGF receptor) (EC 2.7.10.1) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), METImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Storage Buffer	
ClonalityPolyclonalAliasHepatocyte growth factor receptor (HGF receptor) (EC 2.7.10.1) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), METImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Purification Method	>95%, Protein G purified
AliasHepatocyte growth factor receptor (HGF receptor) (EC 2.7.10.1) (HGF/SF receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), METImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Isotype	IgG
receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor) (Tyrosine-protein kinase Met), MET     Immunogen Species   Homo sapiens (Human)     Research Area   Signal Transduction	Clonality	Polyclonal
Research Area Signal Transduction	Alias	receptor) (Proto-oncogene c-Met) (Scatter factor receptor) (SF receptor)
	Immunogen Species	Homo sapiens (Human)
Target Names MET	Research Area	Signal Transduction
	Target Names	MET

1