






## NGFR Antibody

<b>Product Code</b>	CSB-PA10479A0Rb
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P08138
Immunogen	Recombinant Human Tumor necrosis factor receptor superfamily member 16 protein (277-391AA)
Raised In	Rabbit
Species Reactivity	Human, Mouse
<b>Tested Applications</b>	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
<b>Purification Method</b>	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Tumor necrosis factor receptor superfamily member 16 (Gp80-LNGFR) (Low affinity neurotrophin receptor p75NTR) (Low-affinity nerve growth factor receptor) (NGF receptor) (p75 ICD) (CD antigen CD271), NGFR, TNFRSF16
Immunogen Species	Homo sapiens (Human)
Research Area	Neuroscience
Target Names	NGFR





Western Blot

Positive WB detected in: Mouse liver tissue All lanes: NGFR antibody at 2.7µg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 46, 36 kDa Observed band size: 46 kDa




## **CUSABIO TECHNOLOGY LLC**











IHC image of CSB-PA10479A0Rb diluted at 1:400 and staining in paraffin-embedded human glioma performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.