

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

IDH1 Antibody, HRP conjugated

| Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonal | | |
|--|---------------------|---|
| Uniprot No.O75874ImmunogenRecombinant Human Isocitrate dehydrogenase [NADP] cytoplasmic prote (1-240AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasIsocitrate dehydrogenase [NADP] cytoplasmic (IDH) (EC 1.1.1.42) (Cytos NADP-isocitrate dehydrogenase) (IDP) (NADP(+)-specific ICDH) (Oxalosuccinate decarboxylase), IDH1, PICD | Product Code | CSB-PA010989EB01HU |
| ImmunogenRecombinant Human Isocitrate dehydrogenase [NADP] cytoplasmic protect (1-240AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasIsocitrate dehydrogenase [NADP] cytoplasmic (IDH) (EC 1.1.1.42) (Cytos NADP-isocitrate dehydrogenase) (IDP) (NADP(+)-specific ICDH) (Oxalosuccinate decarboxylase), IDH1, PICD | Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Image: Construct of the second seco | Uniprot No. | O75874 |
| Species ReactivityHumanTested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasIsocitrate dehydrogenase [NADP] cytoplasmic (IDH) (EC 1.1.1.42) (Cytos NADP-isocitrate dehydrogenase) (IDP) (NADP(+)-specific ICDH) (Oxalosuccinate decarboxylase), IDH1, PICD | Immunogen | Recombinant Human Isocitrate dehydrogenase [NADP] cytoplasmic protein (1-240AA) |
| Tested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasIsocitrate dehydrogenase [NADP] cytoplasmic (IDH) (EC 1.1.1.42) (Cytos NADP-isocitrate dehydrogenase), IDH1, PICD | Raised In | Rabbit |
| FormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasIsocitrate dehydrogenase [NADP] cytoplasmic (IDH) (EC 1.1.1.42) (Cytos NADP-isocitrate dehydrogenase), IDP) (NADP(+)-specific ICDH) (Oxalosuccinate decarboxylase), IDH1, PICD | Species Reactivity | Human |
| ConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasIsocitrate dehydrogenase [NADP] cytoplasmic (IDH) (EC 1.1.1.42) (Cytos NADP-isocitrate dehydrogenase) (IDP) (NADP(+)-specific ICDH) (Oxalosuccinate decarboxylase), IDH1, PICD | Tested Applications | ELISA |
| Storage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasIsocitrate dehydrogenase [NADP] cytoplasmic (IDH) (EC 1.1.1.42) (Cytos NADP-isocitrate dehydrogenase) (IDP) (NADP(+)-specific ICDH) (Oxalosuccinate decarboxylase), IDH1, PICD | Form | Liquid |
| Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasIsocitrate dehydrogenase [NADP] cytoplasmic (IDH) (EC 1.1.1.42) (Cytos NADP-isocitrate dehydrogenase) (IDP) (NADP(+)-specific ICDH) (Oxalosuccinate decarboxylase), IDH1, PICD | Conjugate | HRP |
| Isotype IgG Clonality Polyclonal Alias Isocitrate dehydrogenase [NADP] cytoplasmic (IDH) (EC 1.1.1.42) (Cytos NADP-isocitrate dehydrogenase) (IDP) (NADP(+)-specific ICDH) (Oxalosuccinate decarboxylase), IDH1, PICD | Storage Buffer | |
| Clonality Polyclonal Alias Isocitrate dehydrogenase [NADP] cytoplasmic (IDH) (EC 1.1.1.42) (Cytos NADP-isocitrate dehydrogenase) (IDP) (NADP(+)-specific ICDH) (Oxalosuccinate decarboxylase), IDH1, PICD | Purification Method | >95%, Protein G purified |
| Alias Isocitrate dehydrogenase [NADP] cytoplasmic (IDH) (EC 1.1.1.42) (Cytos NADP-isocitrate dehydrogenase) (IDP) (NADP(+)-specific ICDH) (Oxalosuccinate decarboxylase), IDH1, PICD | Isotype | IgG |
| NADP-isocitrate dehydrogenase) (IDP) (NADP(+)-specific ICDH) (Oxalosuccinate decarboxylase), IDH1, PICD | Clonality | Polyclonal |
| Immunogen Species Homo sapiens (Human) | Alias | |
| | Immunogen Species | Homo sapiens (Human) |
| Research Area Signal Transduction | Research Area | Signal Transduction |
| Target Names IDH1 | Target Names | IDH1 |

1