



# Recombinant Rat M-phase inducer phosphatase 2 (Cdc25b)

<b>Product Code</b>	CSB-YP004995RA
<b>Storage</b>	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
<b>Uniprot No.</b>	P48966
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Rattus norvegicus (Rat)
<b>Purity</b>	>85% (SDS-PAGE)
<b>Sequence</b>	MEVPPQKSAP GSALSTARVL GGIQRPRHLS GFGFGSDGLL GSPERAASSS PVTTLTQTM Y NLAGLGSETP KTQVGSLSFQ NRLTDLSLSR RTSECSLSSE SSESSDAGLC MDSPSPMDPQ TAERTFEQAI QAASRVIQKM QFTIKASVFA SEAAGHSPVL QNITNSQALD SWEKDEAGYR AASSPGEDKE NDGYIFKMPQ KLPHSSSARA LAEWASRREA FTQRPSSAPD LMCLTTDGKM DVEEASPVAQ SSSLTPVERA CEEDDGFVDI LESDLKDDDM VPAGMENLIS APLVKKLDKE EEQDLIMFSK CQRLFRSPSM PCSVIRPILK RLERPHDRDV PVLKRRKSG TPLEEQLEE PKARVFRSRS LCHEIESILD SDHRGLIGDY SKAFLLQTVD GKHQDLKYIS PETMVALLTG KFSNIVEKFV IVDCRYPYEY EGGHIKNAVN LPLEPDAETF LLKHPITPCN LDKRIILIFH CEFSSERGPR MCRFIRERDR AANDYPSLYY PEMYILKGGY KEFFPQHNF CEPQDYRPMN HAAFRDELRN FRLKTRSWAG ERSTTQLCSR LQDQ
<b>Source</b>	Yeast
<b>Target Names</b>	Cdc25b
<b>Protein Names</b>	Recommended name: M-phase inducer phosphatase 2 EC= 3.1.3.48 Alternative name(s): Dual specificity phosphatase Cdc25B
<b>Expression Region</b>	1-574
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	Full length protein
<b>Target Details</b>	CDC25B is a member of the CDC25 family of phosphatases. CDC25B activates the cyclin dependent kinase CDC2 by removing two phosphate groups and it is required for entry into mitosis. CDC25B shuttles between the nucleus and the cytoplasm due to nuclear localization and nuclear export signals. The protein is nuclear in the M and G1 phases of the cell cycle and moves to the cytoplasm during S and G2. CDC25B has oncogenic properties, although its role in tumor formation has not been determined. Multiple transcript variants for this gene exist.
<b>Reconstitution</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a



concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

### Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.