



# Recombinant Human Cyclin-dependent kinase 8 (CDK8)

<b>Product Code</b>	CSB-YP005077HU
<b>Storage</b>	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
<b>Uniprot No.</b>	P49336
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Purity</b>	>85% (SDS-PAGE)
<b>Sequence</b>	MDYDFKVKLS SERERVEDLF EYEGCKVGRG TYGHVYKAKR KDGKDDKDYA LKQIEGTGIS MSACREIALL RELKHPNVIS LQKVFLSHAD RKVWLLFDYA EHDLWHIIKF HRASKANKKP VQLPRGMVKS LLYQILDGIH YLHANWVLHR DLKPANILVM GEGPERGRVK IADMGFARLF NSPLKPLADL DPVVVTFWYR APELLLGARH YTKAIDIWAI GCIFAELLTS EPIFHCRQED IKTSNPYHHD QLDRIFNVMG FPADKDWEDI KKMPEHSTLM KDFRRNTYTN CSLIKYMEKH KVKPDSKAFH LLQKLLTMDP IKRITSEQAM QDPYFLEDPL PTSDVFAGCQ IPYPKREFLT EEEPDDKGDK KNQQQQQGNN HTNGTGHPGN QDSSHTQGPP LKKVRVVPPT TTSGLIMITS DYQRSNPHAA YPNPGPSTSQ PQSSMGYSAT SQQPPQYSHQ THRY
<b>Source</b>	Yeast
<b>Target Names</b>	CDK8
<b>Protein Names</b>	Recommended name: Cyclin-dependent kinase 8 EC= 2.7.11.22 EC= 2.7.11.23 Alternative name(s): Cell division protein kinase 8 Mediator complex subunit CDK8 Mediator of RNA polymerase II transcription subunit CDK8 Protein ki
<b>Expression Region</b>	1-464
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	Full length protein
<b>Target Details</b>	This protein is a member of the cyclin-dependent protein kinase (CDK) family. CDK family members are highly similar to the gene products of <i>Saccharomyces cerevisiae</i> cdc28, and <i>Schizosaccharomyces pombe</i> cdc2, and are known to be important regulators of cell cycle progression. This kinase and its regulatory subunit cyclin C are components of the RNA polymerase II holoenzyme complex, which phosphorylates the carboxy-terminal domain (CTD) of the largest subunit of RNA polymerase II. This kinase has also been shown to regulate transcription by targeting the CDK7/cyclin H subunits of the general transcription initiation factor IIH (TFIIH), thus providing a link between the Mediator-like protein complexes and the basal transcription machinery.
<b>Reconstitution</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the



contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

---

**Shelf Life**

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.