




NTSR2 Antibody

| | |
|----------------------------|---|
| Product Code | CSB-PA016138LA01HU |
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | O95665 |
| Immunogen | Recombinant Human Neurotensin receptor type 2 protein (359-410AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200, IF:1:50-1:200 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | Neurotensin receptor type 2 (NT-R-2) (NTR2) (Levocabastine-sensitive neurotensin receptor), NTSR2 |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Neuroscience |
| Target Names | NTSR2 |

Image



Western Blot

Positive WB detected in: U87 whole cell lysate


All lanes: NTSR2 antibody at 3.2µg/ml

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 46 kDa


Observed band size: 46 kDa



Immunohistochemistry of paraffin-embedded human adrenal gland tissue using CSB-PA016138LA01HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human prostate tissue using CSB-PA016138LA01HU at dilution of 1:100



Immunofluorescence staining of HeLa cells with CSB-PA016138LA01HU at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).