



🕜 Tel: +1-301-363-4651 🛛 🗵 Email: cusabio@cusabio.com 📀 Website: www.cusabio.com 🍙

## **BCAM** Antibody

Product Code	CSB-PA341920LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P50895
Immunogen	Recombinant Human Basal cell adhesion molecule protein (447-547AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:200-1:500
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Basal cell adhesion molecule (Auberger B antigen) (B-CAM cell surface glycoprotein) (F8/G253 antigen) (Lutheran antigen) (Lutheran blood group glycoprotein) (CD antigen CD239), BCAM, LU MSK19
Immunogen Species	Homo sapiens (Human)
Research Area	Cardiovascular
Target Names	BCAM
Image	Immunohistochemistry of poroffin omhoddod



Immunohistochemistry of paraffin-embedded human pancreatic cancer using CSB-PA341920LA01HU at dilution of 1:100




Immunohistochemistry of paraffin-embedded human kidney tissue using CSB-PA341920LA01HU at dilution of 1:100

1



## **CUSABIO TECHNOLOGY LLC**

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 📀 Website: www.cusabio.com 🍵



Immunofluorescence staining of HepG2 cells with CSB-PA341920LA01HU at 1:200, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).