




USP10 Antibody

Product Code	CSB-PA613516LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q14694
Immunogen	Recombinant Human Ubiquitin carboxyl-terminal hydrolase 10 protein (453-592AA)
Raised In	Rabbit
Species Reactivity	Human, Rat
Tested Applications	ELISA, WB, IHC, IF, IP; Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200, IF:1:50-1:200, IP:1:200-1:2000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Ubiquitin carboxyl-terminal hydrolase 10 (EC 3.4.19.12) (Deubiquitinating enzyme 10) (Ubiquitin thioesterase 10) (Ubiquitin-specific-processing protease 10), USP10, KIAA0190
Immunogen Species	Homo sapiens (Human)
Research Area	Cell Biology
Target Names	USP10

Image



Western Blot

Positive WB detected in: HeLa whole cell lysate, 293 whole cell lysate, A549 whole cell lysate, Rat liver tissue


All lanes: USP10 antibody at 3µg/ml

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 88, 93 kDa


Observed band size: 88 kDa



Immunohistochemistry of paraffin-embedded human testis tissue using CSB-PA613516LA01HU at dilution of 1:100



Immunofluorescent analysis of HeLa cells using CSB-PA613516LA01HU at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Immunoprecipitating USP10 in A549 whole cell lysate
 Lane 1: Rabbit control IgG instead of CSB-PA613516LA01HU in A549 whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)
 Lane 2: CSB-PA613516LA01HU (8μg) + A549 whole cell lysate (500μg)
 Lane 3: A549 whole cell lysate (10μg)

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.