




NEDD4L Antibody

| | |
|----------------------------|---|
| Product Code | CSB-PA850411LA01HU |
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q96PU5 |
| Immunogen | Recombinant Human E3 ubiquitin-protein ligase NEDD4-like protein (289-494AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, WB, IF; Recommended dilution: WB:1:500-1:5000, IF:1:50-1:200 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | E3 ubiquitin-protein ligase NEDD4-like (EC 2.3.2.26) (HECT-type E3 ubiquitin transferase NED4L) (NEDD4.2) (Nedd4-2), NEDD4L, KIAA0439 NEDL3 |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Epigenetics and Nuclear Signaling |
| Target Names | NEDD4L |

Image




Western Blot

Positive WB detected in: 293 whole cell lysate, Jurkat whole cell lysate, HeLa whole cell lysate
All lanes: NEDD4L antibody at 3µg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution
Predicted band size: 112, 105, 101, 99, 111, 110, 97 kDa
Observed band size: 112 kDa



Immunofluorescence staining of HepG2 cells with CSB-PA850411LA01HU at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.