


🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 📀 Website: www.cusabio.com 🌘

## MAP3K4 Antibody

| Product Code               | CSB-PA013426LA01HU   |
|----------------------------|--|
| Storage                    | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  |
| Uniprot No.                | Q9Y6R4   |
| Immunogen                  | Recombinant Human Mitogen-activated protein kinase kinase kinase 4 protein (53-145AA)  |
| Raised In                  | Rabbit   |
| Species Reactivity         | Human  |
| <b>Tested Applications</b> | ELISA, IHC; Recommended dilution: IHC:1:20-1:200   |
| Form                       | Liquid   |
| Conjugate                  | Non-conjugated   |
| Storage Buffer             | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, pH 7.4   |
| Purification Method        | >95%, Protein G purified   |
| Isotype                    | lgG  |
| Clonality                  | Polyclonal   |
| Alias                      | Mitogen-activated protein kinase kinase kinase 4 (EC 2.7.11.25) (MAP three kinase 1) (MAPK/ERK kinase kinase 4) (MEK kinase 4) (MEKK 4), MAP3K4, KIAA0213 MAPKKK4 MEKK4 MTK1 |
| Immunogen Species          | Homo sapiens (Human)   |
| Research Area              | Signal Transduction  |
| Target Names               | MAP3K4   |
|                            |  |

Image




IHC image of CSB-PA013426LA01HU diluted at 1:100 and staining in paraffin-embedded human pancreatic tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

1



🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 📀 Website: www.cusabio.com 🍵



IHC image of CSB-PA013426LA01HU diluted at 1:100 and staining in paraffin-embedded human heart tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated ABC system.