






UIMC1 Antibody

Product Code	CSB-PA857019LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q96RL1
Immunogen	Recombinant Human BRCA1-A complex subunit RAP80 protein (161-235AA)
Raised In	Rabbit
Species Reactivity	Human, Rat
Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:500-1:1000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	BRCA1-A complex subunit RAP80 (Receptor-associated protein 80) (Retinoid X receptor-interacting protein 110) (Ubiquitin interaction motif-containing protein 1), UIMC1, RAP80 RXRIP110
Immunogen Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Target Names	UIMC1
Image	Western Rlot



Western Blot

Positive WB detected in: HepG2 whole cell lysate, A549 whole cell lysate, Rat kidney tissue

All lanes: UIMC1 antibody at 3µg/ml

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 80, 62, 71, 39, 18 kDa

Observed band size: 80 kDa




CUSABIO TECHNOLOGY LLC









IHC image of CSB-PA857019LA01HU diluted at 1:500 and staining in paraffin-embedded human testis tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.