


🕜 Tel: +1-301-363-4651 🛛 🖾 Email: cusabio@cusabio.com 🛛 🥑 Website: www.cusabio.com 🌘


GP6 Antibody

Product Code	CSB-PA872550LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9HCN6
Immunogen	Recombinant Human Platelet glycoprotein VI protein (115-265AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, IHC, IF; Recommended dilution: IHC:1:500-1:1000, IF:1:200-1:500
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
lsotype	IgG
Clonality	Polyclonal
Alias	Platelet glycoprotein VI (GPVI) (Glycoprotein 6), GP6
Immunogen Species	Homo sapiens (Human)
Research Area	Cardiovascular
Target Names	GP6

Image



IHC image of CSB-PA872550LA01HU diluted at 1:600 and staining in paraffin-embedded human bladder cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of CSB-PA872550LA01HU diluted at 1:600 and staining in paraffin-embedded human glioma performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.


1



CUSABIO TECHNOLOGY LLC

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 📀 Website: www.cusabio.com 🍵





Immunofluorescence staining of MCF-7 cells with CSB-PA872550LA01HU at 1:200, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).