


🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🛛 🥭 Website: www.cusabio.com 🌘


PLXNB1 Antibody

Product Code	CSB-PA018222LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	O43157
Immunogen	Recombinant Human Plexin-B1 protein (664-837AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, IHC, IF; Recommended dilution: IHC:1:200-1:500, IF:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
lsotype	lgG
Clonality	Polyclonal
Alias	Plexin-B1 (Semaphorin receptor SEP), PLXNB1, KIAA0407 PLXN5 SEP
Immunogen Species	Homo sapiens (Human)
Research Area	Neuroscience
Target Names	PLXNB1

Image



IHC image of CSB-PA018222LA01HU diluted at 1:200 and staining in paraffin-embedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.




IHC image of CSB-PA018222LA01HU diluted at 1:200 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

1



🕜 Tel: +1-301-363-4651 🛛 🗵 Email: cusabio@cusabio.com 📀 Website: www.cusabio.com 🍙



Immunofluorescence staining of Hela cells with CSB-PA018222LA01HU at 1:66, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).