






HIST1H3A (Ab-36) Antibody

Product Code Storage Uniprot No. Immunogen	CSB-PA010418PA36nme1HU Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. P68431
Uniprot No.	
· · · · · · · · · · · · · · · · · · ·	P68431
Immunogen	
	Peptide sequence around site of Lys (36) derived from Human Histone H3.1
Raised In	Rabbit
Species Reactivity	Human, Mouse
Tested Applications	ELISA, WB, IHC, IF, ChIP; Recommended dilution: WB:1:200-1:2000, IHC:1:20-1:200, IF:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	Antigen Affinity Purified
Isotype	IgG
Clonality	Polyclonal
Alias	Histone H3.1 (Histone H3/a) (Histone H3/b) (Histone H3/c) (Histone H3/d) (Histone H3/f) (Histone H3/h) (Histone H3/i) (Histone H3/j) (Histone H3/k) (Histone H3/l), HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J, H3FA; H3FL; H3FC; H3FB; H3FD; H3FI; H3FH; H3FK; H3FF; H3FJ
Immunogen Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Target Names	HIST1H3A
Image	



Western Blot

Positive WB detected in: Hela cell acid extracts, HEK293 cell acid extracts, Mouse liver tissue, Mouse kidney tissue

All lanes: HIST1H3A antibody at 2µg/ml


Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 16 kDa Observed band size: 16 kDa

















Immunohistochemistry of paraffin-embedded human colon cancer using CSB-PA010418PA36nme1HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human lung cancer using CSB-PA010418PA36nme1HU at dilution of 1:100



Immunofluorescent analysis of Hela cells using CSB-PA010418PA36nme1HU at dilution of 1:100 and Cy3-congugated Goat Anti-Rabbit IgG



Chromatin Immunoprecipitation Hela (4*10⁶) were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with 8µg anti-HIST1H3A (CSB-PA010418PA36nme1HU) or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers against the β -Globin promoter.