







## Acetyl-HIST1H1E (K16) Antibody

<b>Product Code</b>	CSB-PA010380PA16acHU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P10412
Immunogen	Peptide sequence around site of Acetyl-Lys (16) derived from Human Histone H1.4
Raised In	Rabbit
Species Reactivity	Human
<b>Tested Applications</b>	ELISA, ICC, IF, ChIP; Recommended dilution: ICC:1:20-1:200, IF:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	Antigen Affinity Purified
Isotype	IgG
Clonality	Polyclonal
Alias	Histone H1.4 (Histone H1b) (Histone H1s-4), HIST1H1E, H1F4
Immunogen Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Target Names	HIST1H1E
Image	Immunocytochomistry analysis of MCE 7 colls



Immunocytochemistry analysis of MCF-7 cells using CSB-PA010380PA16acHU at dilution of 1:100



Immunofluorescent analysis of MCF-7 cells (sodium butyrate, 30 mM, 4h) using CSB-PA010380PA16acHU at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)




## **CUSABIO TECHNOLOGY LLC**



🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 🕒 Website: www.cusabio.com 🥑







Chromatin Immunoprecipitation Hela (4\*106, treated with 30mM sodium butyrate for 4h) were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with 8µg anti-HIST1H1E (CSB-PA010380PA16acHU) or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers against the  $\beta$ -Globin promoter.