




# Mono-methyl-H1F0 (K11) Antibody

<b>Product Code</b>	CSB-PA010087PA11me1HU
<b>Abbreviation</b>	Histone H1.0
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	P07305
<b>Immunogen</b>	Peptide sequence around site of Mono-methyl-Lys (11) derived from Human Histone H1.0
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, IF; Recommended dilution: IF:1:50-1:200
<b>Relevance</b>	Histones H1 are necessary for the condensation of nucleosome chains into higher-order structures. The H1F0 histones are found in cells that are in terminal stages of differentiation or that have low rates of cell division.
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
<b>Purification Method</b>	Antigen Affinity Purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Histone H1.0 (Histone H1') (Histone H1(0)) [Cleaved into: Histone H1.0, N-terminally processed], H1F0, H1FV
<b>Species</b>	Human
<b>Research Area</b>	Epigenetics and Nuclear Signaling
<b>Target Names</b>	H1F0

## Image



Immunofluorescent analysis of HepG2 cells using CSB-PA010087PA11me1HU at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)