




Acetyl-Histone H2B type 1-B (K20) Recombinant Monoclonal Antibody

Product Code	CSB-RA010402A20acHU
Abbreviation	Histone H2B type 1-B
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P33778
Immunogen	A synthesized peptide
Species Reactivity	Human
Tested Applications	ELISA, WB, ICC, IF, FC; Recommended dilution: WB:1:5000-1:10000, ICC:1:50-1:500, IF:1:30-1:200
Relevance	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Purification Method	Affinity-chromatography
Isotype	Rabbit IgG
Clonality	Monoclonal
Product Type	Recombinant Antibody
Immunogen Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Gene Names	HIST1H2BB
Clone No.	21F11
Image	



Western Blot

Positive WB detected in HeLa whole cell lysate treated by 15mM sodium butyrate for 30min


All lanes Acetyl-Histone H2B type 1-B(K20)antibody at 0.135μg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution


Predicted band size: 15 KDa

Observed band size: 15 KDa




Immunocytochemistry analysis of CSB-

RA010402A20acHU diluted at 1:100 and staining in HeLa cells performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of HeLa cells (treated by 15mM sodium butyrate for 30min) with CSB-RA010402A20acHU at 1:84, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



Overlay histogram showing HeLa cells stained with CSB-RA010402A20acHU (red line) at 1:50. The cells were fixed with 70% Ethylalcohol (18h) and then permeabilized with 0.3% Triton X-100 for 2 min. The cells were then incubated in 1x PBS /10% normal goat serum to block non-specific protein-protein interactions followed by primary antibody for 1 h at 4°C. The secondary antibody used was FITC goat anti-rabbit IgG (H+L) at 1/200 dilution for 1 h at 4°C. Control antibody (green line) was used under the same conditions. Acquisition of >10,000 events was performed.

Description

The acetyl-Histone H2B type 1-B/HIST1H2BB (K20) recombinant monoclonal antibody is a highly specific antibody against the acetylated HIST1H2BB at K20 from human and mouse sources. This acetyl-HIST1H2BB (K20) antibody was expressed by transfecting the human acetyl-HIST1H2BB (K20) monoclonal antibody gene-vector clones into the cell line for in vitro production and



subsequent purification from the tissue culture supernatant (TCS) through affinity-chromatography. Its isotype matches with the rabbit IgG. This acetyl-HIST1H2BB (K20) antibody can be used in ELISA, WB, ICC, IF, and FC applications.

HIST1H2BB is a subtype of histone core component H2B. Antibodies against the HIST1H2BB are useful for the localization and detection of the HIST1H2BB of acetylation at Lys20 residue.