

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

DLST Antibody, HRP conjugated

StorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.P36957ImmunogenRecombinant Human Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial protein (76-225AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasDihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.		
Uniprot No.P36957ImmunogenRecombinant Human Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial protein (76-225AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasDihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate <br< td=""><td>Product Code</td><td>CSB-PA861526LB01HU</td></br<>	Product Code	CSB-PA861526LB01HU
ImmunogenRecombinant Human Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial protein (76-225AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasDihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex, component E2) (OGDC-E2) (Dihydrolipoamide succinyltransferase component of 2-oxoglutarate dehydrogenase complex, component E2) (OGDC-E2) (Dihydrolipoamide succinyltransferase component of 2-oxoglutarate dehydrogenase complex component E3) (OGDC-E2) (Dihydrolipoamide succinyltransferase component of 2-oxoglutarate dehydrogenase complex component E3) (OGDC-E2) (Dihydrolipoamide succin	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
component of 2-oxoglutarate dehydrogenase complex, mitochondrial protein (76-225AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasDihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex component of 2-oxoglutarate dehydrogenase complex mitochondrial (EC 2.3.1.61) (E2K), DLST, DLTSImmunogen SpeciesHomo sapiens (Human)	Uniprot No.	P36957
Species ReactivityHumanTested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasDihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex component f2) (OGDC-E2) (Dihydrolipoamide succinyltransferase complex to 2-oxoglutarate dehydrogenase complex) (E2K), DLST, DLTSImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Immunogen	component of 2-oxoglutarate dehydrogenase complex, mitochondrial protein
Tested ApplicationsELISAFormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasDihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex component E2) (OGDC-E2) (Dihydrolipoamide succinyltransferase complex) (E2K), DLST, DLTSImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Raised In	Rabbit
FormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasDihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex component E2) (OGDC-E2) (Dihydrolipoamide succinyltransferase complex) (E2K), DLST, DLTSImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Species Reactivity	Human
ConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasDihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex, motochondrial (EC 2.3.1.61) (2-oxoglu	Tested Applications	ELISA
Storage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasDihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex component E2) (OGDC-E2) (Dihydrolipoamide succinyltransferase complex component of 2-oxoglutarate dehydrogenase complex component of 2-oxoglutarate dehydrogenase complex) (E2K), DLST, DLTSImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Form	Liquid
Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasDihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex component E2) (OGDC-E2) (Dihydrolipoamide succinyltransferase complex of 2-oxoglutarate dehydrogenase complex component of 2-oxoglutarate dehydrogenase complex component E2) (OGDC-E2) (Dihydrolipoamide succinyltransferase component of 2-oxoglutarate dehydrogenase complex) (E2K), DLST, DLTSImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Conjugate	HRP
IsotypeIgGClonalityPolyclonalAliasDihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex component E2) (OGDC-E2) (Dihydrolipoamide succinyltransferase component of 2-oxoglutarate dehydrogenase complex) (E2K), DLST, DLTSImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Storage Buffer	
ClonalityPolyclonalAliasDihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrolipoamide succinyltransferase component of 2-oxoglutarate dehydrogenase complex (E2K), DLST, DLTSImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Purification Method	>95%, Protein G purified
AliasDihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex component E2) (OGDC-E2) (Dihydrolipoamide succinyltransferase component of 2-oxoglutarate dehydrogenase complex) (E2K), DLST, DLTSImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Isotype	IgG
dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex component E2) (OGDC-E2) (Dihydrolipoamide succinyltransferase component of 2-oxoglutarate dehydrogenase complex) (E2K), DLST, DLTSImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Clonality	Polyclonal
Research Area Signal Transduction	Alias	dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex component E2) (OGDC-E2) (Dihydrolipoamide succinyltransferase component of 2-oxoglutarate dehydrogenase complex)
	Immunogen Species	Homo sapiens (Human)
Target Names DLST	Research Area	Signal Transduction
	Target Names	DLST

1