




TNFRSF19 Antibody

| Product Code | CSB-PA878878LA01HU |
|----------------------------|---|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q9NS68 |
| Immunogen | Recombinant Human Tumor necrosis factor receptor superfamily member 19 protein (276-415AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, WB, IF; Recommended dilution: WB:1:500-1:5000, IF:1:50-1:200 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | Tumor necrosis factor receptor superfamily member 19 (TRADE) (Toxicity and JNK inducer), TNFRSF19, TAJ TROY |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Cell Biology |
| Target Names | TNFRSF19 |
| Image | wastern Blot |



Positive WB detected in: PC-3 whole cell lysate All lanes: TNFRSF19 antibody at 2.36μg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 47, 46, 31 kDa Observed band size: 47 kDa




CUSABIO TECHNOLOGY LLC











Immunofluorescence staining of HepG2 cells with CSB-PA878878LA01HU at 1:200, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).