





# Acetyl-HIST1H3A (K4) Antibody

<b>Product Code</b>	CSB-PA010418OA04acHU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	P68431
<b>Immunogen</b>	Peptide sequence around site of Acetyl-Lys (4) derived from Human Histone H3.1
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IF, IP, ChIP; Recommended dilution: WB:1:200-1:2000, IF:1:1-1:10, IP:1:200-1:2000
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
<b>Purification Method</b>	Antigen Affinity Purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Histone H3.1 (Histone H3/a) (Histone H3/b) (Histone H3/c) (Histone H3/d) (Histone H3/f) (Histone H3/h) (Histone H3/i) (Histone H3/j) (Histone H3/k) (Histone H3/l), HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J, H3FA; H3FL; H3FC; H3FB; H3FD; H3FI; H3FH; H3FK; H3FF; H3FJ
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Epigenetics and Nuclear Signaling
<b>Target Names</b>	HIST1H3A


## Image




Immunofluorescence staining of HeLa cells with CSB-PA010418OA04acHU at 1:7.5, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Chromatin Immunoprecipitation HeLa ( $4 \times 10^6$ , treated with 30mM sodium butyrate for 4h) were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with 5µg anti-HIST1H3A (CSB-PA010418OA04acHU) or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers against the  $\beta$ -Globin promoter.



**Western Blot**  
 Positive WB detected in: Jurkat whole cell lysate, 293 whole cell lysate, HepG2 whole cell lysate (treated by 30mM sodium butyrate for 4h)  
 All lanes: HIST1H3A antibody at 0.935µg/ml  
 Secondary  
 Goat polyclonal to rabbit IgG at 1/50000 dilution  
 Predicted band size: 16 kDa  
 Observed band size: 16 kDa



Immunoprecipitating HIST1H3A in HepG2 whole cell lysate (treated with 30mM sodium butyrate for 4h)  
 Lane 1: Rabbit control IgG instead of CSB-PA010418OA04acHU in HepG2 whole cell lysate (treated with 30mM sodium butyrate for 4h). For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)  
 Lane 2: CSB-PA010418OA04acHU (5µg) + HepG2 whole cell lysate (treated with 30mM sodium butyrate for 4h) (500µg)  
 Lane 3: HepG2 whole cell lysate (treated with 30mM sodium butyrate for 4h) (20µg)

**Usage**

For Research Use Only. Not for use in diagnostic or therapeutic procedures.