









## MFAP4 Antibody

| <b>Product Code</b>        | CSB-PA013749LA01HU   |
|----------------------------|--|
| Storage                    | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.                    |
| Uniprot No.                | P55083   |
| Immunogen                  | Recombinant Human Microfibril-associated glycoprotein 4 protein (100-191AA)      |
| Raised In                  | Rabbit   |
| Species Reactivity         | Human  |
| <b>Tested Applications</b> | ELISA, IHC; Recommended dilution: IHC:1:200-1:500                                |
| Form                       | Liquid   |
| Conjugate                  | Non-conjugated   |
| Storage Buffer             | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Purification Method        | >95%, Protein G purified   |
| Isotype                    | IgG  |
| Clonality                  | Polyclonal   |
| Alias                      | Microfibril-associated glycoprotein 4, MFAP4                                     |
| Immunogen Species          | Homo sapiens (Human)   |
| Research Area              | Signal Transduction  |
| Target Names               | MFAP4  |
| Image                      | ILIC image of CCD DA013740LA01LILI diluted of                                    |



IHC image of CSB-PA013749LA01HU diluted at 1:300 and staining in paraffin-embedded human prostate cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of CSB-PA013749LA01HU diluted at 1:300 and staining in paraffin-embedded human cervical cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



## **CUSABIO TECHNOLOGY LLC**



