




Propionyl-HIST1H4A (K8) Antibody

Product Code	CSB-PA010429OA08propHU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P62805
Immunogen	Peptide sequence around site of Propionyl-Lys (8) derived from Human Histone H4
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IF, IP, ChIP; Recommended dilution: WB:1:1000-1:5000, IF:1:20-1:200, IP:1:200-1:2000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	Antigen Affinity Purified
Isotype	IgG
Clonality	Polyclonal
Alias	Histone H4, HIST1H4A; HIST1H4B; HIST1H4C; HIST1H4D; HIST1H4E; HIST1H4F; HIST1H4H; HIST1H4I; HIST1H4J; HIST1H4K; HIST1H4L; HIST2H4A; HIST2H4B; HIST4H4, H4/A H4FA; H4/I H4FI; H4/G H4FG; H4/B H4FB; H4/J H4FJ; H4/C H4FC; H4/H H4FH; H4/M H4FM; H4/E H4FE; H4/D H4FD; H4/K H4FK; H4/N H4F2 H4FN HIST2H4; H4/O H4FO;
Immunogen Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Target Names	HIST1H4A

Image




Western Blot


Detected samples: Hela whole cell lysate, A549 whole cell lysate, 293 whole cell lysate, K562 whole cell lysate; Untreated (-) or treated (+) with 10mM sodium propionate for 4h
All lanes: HIST1H4A antibody at 1:2000

Secondary

Goat polyclonal to rabbit IgG at 1/40000 dilution
Predicted band size: 12 kDa
Observed band size: 12 kDa



Immunofluorescence staining of HeLa cells (treated with 10mM sodium propionate for 4h) with CSB-PA010429OA08propHU at 1:25, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




Immunoprecipitating H2A in HeLa whole cell lysate

Lane 1: Rabbit control IgG instead of CSB-PA010429OA08propHU in HeLa whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)

Lane 2: CSB-PA010429OA08propHU (5μg) + HeLa whole cell lysate (500μg)

Lane 3: HeLa whole cell lysate (20μg)



Chromatin Immunoprecipitation HeLa (4×10^6 , treated with 10mM sodium propionate for 4h) were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with 5μg anti-H2A (CSB-PA010429OA08propHU) or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers against the β -Globin promoter.