






HIST1H2BC (Ab-116) Antibody

Product Code	CSB-PA010403OA116nme1HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P62807
Immunogen	Peptide sequence around site of Lys (116) derived from Human Histone H2B type 1-C/E/F/G/I
Raised In	Rabbit
Species Reactivity	Human, Mouse
Tested Applications	ELISA, WB, IP, ChIP; Recommended dilution: WB:1:100-1:1000, IP:1:200-1:2000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	Antigen Affinity Purified
Isotype	IgG
Clonality	Polyclonal
Alias	Histone H2B type 1-C/E/F/G/I (Histone H2B.1 A) (Histone H2B.a) (H2B/a) (Histone H2B.g) (H2B/g) (Histone H2B.h) (H2B/h) (Histone H2B.k) (H2B/k) (Histone H2B.I) (H2B/I), HIST1H2BC; HIST1H2BE; HIST1H2BF; HIST1H2BG; HIST1H2BI, H2BFL; H2BFH; H2BFG; H2BFA; H2BFK
Immunogen Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Target Names	HIST1H2BC
Image	
	Western Blot



Western Blot

Positive WB detected in: Hela whole cell lysate, 293 whole cell lysate, HL60 whole cell lysate, HepG2 whole cell lysate, MCF-7 whole cell lysate, Mouse liver tissue

All lanes: HIST1H2BC antibody at 0.92µg/ml

Goat polyclonal to rabbit IgG at 1/50000 dilution


Predicted band size: 14 kDa Observed band size: 14 kDa












Chromatin Immunoprecipitation Hela (4*10⁶) were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with 5µg anti-HIST1H2BC (CSB-PA010403OA116nme1HU) or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers against the β -Globin promoter.



Immunoprecipitating HIST1H2BC in MCF-7 whole cell lysate

Lane 1: Rabbit control IgG instead of CSB-PA010403OA116nme1HU in MCF-7 whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)

Lane 2: CSB-PA010403OA116nme1HU (5µg) +

MCF-7 whole cell lysate (500µg)

Lane 3: MCF-7 whole cell lysate (20µg)