




PARP3 Antibody

Product Code	CSB-PA017467EA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9Y6F1
Immunogen	Recombinant Human Poly [ADP-ribose] polymerase 3 protein (1-240AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Poly [ADP-ribose] polymerase 3 (PARP-3) (hPARP-3) (EC 2.4.2.30) (ADP-ribosyltransferase diphtheria toxin-like 3) (ARTD3) (IRT1) (NAD(+) ADP-ribosyltransferase 3) (ADPRT-3) (Poly[ADP-ribose] synthase 3) (pADPRT-3), PARP3, ADPRT3 ADPRTL3
Immunogen Species	Homo sapiens (Human)
Research Area	Cell Biology
Target Names	PARP3

Image



Western Blot

Positive WB detected in: A549 whole cell lysate, LO2 whole cell lysate


All lanes: PARP3 antibody at 3.5µg/ml

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 61 kDa

Observed band size: 61 kDa



IHC image of CSB-PA017467EA01HU diluted at 1:100 and staining in paraffin-embedded human tonsil tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of CSB-PA017467EA01HU diluted at 1:100 and staining in paraffin-embedded human glioma performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.