

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com 🍙

PPCDC Antibody, Biotin conjugated

StorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.Q96CD2ImmunogenRecombinant Human Phosphopantothenoylcysteine decarboxylase protein (51-183AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISAFormLiquidConjugateBiotinStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPhosphopantothenoylcysteine decarboxylase (PPC-DC) (EC 4.1.1.36) (CoaC), PPCDC, COACImmunogen SpeciesHomo sapiens (Human)		
Uniprot No.Q96CD2ImmunogenRecombinant Human Phosphopantothenoylcysteine decarboxylase protein (51-183AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISAFormLiquidConjugateBiotinStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPhosphopantothenoylcysteine decarboxylase (PPC-DC) (EC 4.1.1.36) (CoaC), PPCDC, COACImmunogen SpeciesHomo sapiens (Human)	Product Code	CSB-PA018433LD01HU
ImmunogenRecombinant Human Phosphopantothenoylcysteine decarboxylase protein (51-183AA)Raised InRabbitRaised InRabbitSpecies ReactivityHumanTested ApplicationsELISAFormLiquidConjugateBiotinStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPhosphopantothenoylcysteine decarboxylase (PPC-DC) (EC 4.1.1.36) (CoaC), PPCDC, COACImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
ConjugateBiotinStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPhosphopantothenoylcysteine decarboxylase (PPC-DC) (EC 4.1.1.36) (CoaC), PPCDC, COACResearch AreaSignal Transduction	Uniprot No.	Q96CD2
Species ReactivityHumanTested ApplicationsELISAFormLiquidConjugateBiotinStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPhosphopantothenoylcysteine decarboxylase (PPC-DC) (EC 4.1.1.36) (CoaC), PPCDC, COACImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Immunogen	
Tested ApplicationsELISAFormLiquidConjugateBiotinStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPhosphopantothenoylcysteine decarboxylase (PPC-DC) (EC 4.1.1.36) (CoaC), PPCDC, COACImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Raised In	Rabbit
FormLiquidConjugateBiotinStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPhosphopantothenoylcysteine decarboxylase (PPC-DC) (EC 4.1.1.36) (CoaC), PPCDC, COACImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Species Reactivity	Human
ConjugateBiotinStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPhosphopantothenoylcysteine decarboxylase (PPC-DC) (EC 4.1.1.36) (CoaC), PPCDC, COACImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Tested Applications	ELISA
Storage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPhosphopantothenoylcysteine decarboxylase (PPC-DC) (EC 4.1.1.36) (CoaC), PPCDC, COACImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Form	Liquid
Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPhosphopantothenoylcysteine decarboxylase (PPC-DC) (EC 4.1.1.36) (CoaC), PPCDC, COACImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Conjugate	Biotin
IsotypeIgGClonalityPolyclonalAliasPhosphopantothenoylcysteine decarboxylase (PPC-DC) (EC 4.1.1.36) (CoaC), PPCDC, COACImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Storage Buffer	
ClonalityPolyclonalAliasPhosphopantothenoylcysteine decarboxylase (PPC-DC) (EC 4.1.1.36) (CoaC), PPCDC, COACImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Purification Method	>95%, Protein G purified
AliasPhosphopantothenoylcysteine decarboxylase (PPC-DC) (EC 4.1.1.36) (CoaC), PPCDC, COACImmunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction	Isotype	IgG
PPCDC, COAC Immunogen Species Homo sapiens (Human) Research Area Signal Transduction	Clonality	Polyclonal
Research Area Signal Transduction	Alias	
•	Immunogen Species	Homo sapiens (Human)
Target Names PPCDC	Research Area	Signal Transduction
	Target Names	PPCDC

1