


🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🛛 🥥 Website: www.cusabio.com 🌘

MAML2 Antibody

Product Code	CSB-PA809028LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q8IZL2
Immunogen	Recombinant Human Mastermind-like protein 2 protein (347-506AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:500-1:1000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	lgG
Clonality	Polyclonal
Alias	Mastermind-like protein 2 (Mam-2), MAML2, KIAA1819
Immunogen Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Target Names	MAML2

Image



Western Blot


Positive WB detected in: Hela whole cell lysate, SH-SY5Y whole cell lysate, A549 whole cell lysate, PC-3 whole cell lysate, COLO320 whole cell lysate All lanes: MAML2 antibody at 2.2µg/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 126 kDa Observed band size: 126 kDa

1




CUSABIO TECHNOLOGY LLC

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🤅 Website: www.cusabio.com 🍯



IHC image of CSB-PA809028LA01HU diluted at 1:800 and staining in paraffin-embedded human prostate tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of CSB-PA809028LA01HU diluted at 1:800 and staining in paraffin-embedded human adrenal gland tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.