


**Image** 



## MFSD4A Antibody

| <b>Product Code</b>        | CSB-PA850797LA01HU  |
|----------------------------|---|
| Storage                    | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.   |
| Uniprot No.                | Q8N468  |
| Immunogen                  | Recombinant Human Major facilitator superfamily domain-containing protein 4A protein (160-220AA)                                      |
| Raised In                  | Rabbit  |
| <b>Species Reactivity</b>  | Human   |
| <b>Tested Applications</b> | ELISA, WB, IF; Recommended dilution: WB:1:500-1:5000, IF:1:50-1:200   |
| Form                       | Liquid  |
| Conjugate                  | Non-conjugated  |
| Storage Buffer             | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, pH 7.4  |
| <b>Purification Method</b> | >95%, Protein G purified  |
| Isotype                    | IgG   |
| Clonality                  | Polyclonal  |
| Alias                      | Major facilitator superfamily domain-containing protein 4A (Major facilitator superfamily domain-containing protein 4), MFSD4A, MFSD4 |
| Immunogen Species          | Homo sapiens (Human)  |
| Research Area              | Signal Transduction   |
| Target Names               | MFSD4A  |
|                            |   |



Western Blot

Positive WB detected in: A549 whole cell lysate,

HepG2 whole cell lysate

All lanes: MFSD4A antibody at 3.38µg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 57, 47 kDa Observed band size: 47 kDa




## **CUSABIO TECHNOLOGY LLC**









Immunofluorescence staining of Hela cells with CSB-PA850797LA01HU at 1:166, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).