





Acetyl-HIST1H2AG (K13) Antibody


| | |
|----------------------------|--|
| Product Code | CSB-PA010389OA13acHU |
| Abbreviation | Histone H2A type 1 |
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | P0C0S8 |
| Immunogen | Peptide sequence around site of Acetyl-Lys (13) derived from Human Histone H2A type 1 |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, WB, IHC, IF, IP; Recommended dilution: WB:1:100-1:1000, IHC:1:1-1:10, IF:1:1-1:10, IP:1:200-1:2000 |
| Relevance | Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Purification Method | Antigen Affinity Purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | Histone H2A type 1 (H2A.1) (Histone H2A/ptI), HIST1H2AG; HIST1H2AI; HIST1H2AK; HIST1H2AL; HIST1H2AM, H2AFP; H2AFC; H2AFD; H2AFI; H2AFN |
| Species | Human |
| Research Area | Epigenetics and Nuclear Signaling |
| Target Names | HIST1H2AG |
| Image | |


Western Blot


Positive WB detected in: HeLa whole cell lysate, 293 whole cell lysate, A549 whole cell lysate, Jurkat whole cell lysate, HepG2 whole cell lysate (all treated with 30mM sodium butyrate for 4h)
 All lanes: HIST1H2AG antibody at 1.25µg/ml
 Secondary
 Goat polyclonal to rabbit IgG at 1/50000 dilution
 Predicted band size: 15 kDa
 Observed band size: 15 kDa



Immunocytochemistry analysis of CSB-PA010389OA13acHU diluted at 1:5 and staining in HeLa cells (treated with 30mM sodium butyrate for 4h) performed on a Leica Bond™ system. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of HeLa cells (treated with 30mM sodium butyrate for 4h) with CSB-PA010389OA13acHU at 1:2.5, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).


Immunoprecipitating HIST1H2AG in 293 whole cell lysate

Lane 1: Rabbit control IgG instead of CSB-PA010389OA13acHU in 293 whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)
 Lane 2: CSB-PA010389OA13acHU (5µg) + 293 whole cell lysate (500µg)
 Lane 3: 293 whole cell lysate (20µg)