




Crotonyl-HIST1H3A (K9) Antibody

Product Code	CSB-PA010418OA09crHU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P68431
Immunogen	Peptide sequence around site of Crotonyl-Lys (9) derived from Human Histone H3.1
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, ICC, IP, ChIP; Recommended dilution: WB:1:100-1:1000, ICC:1:20-1:200, IP:1:200-1:2000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	Antigen Affinity Purified
Isotype	IgG
Clonality	Polyclonal
Alias	Histone H3.1 (Histone H3/a) (Histone H3/b) (Histone H3/c) (Histone H3/d) (Histone H3/f) (Histone H3/h) (Histone H3/i) (Histone H3/j) (Histone H3/k) (Histone H3/l), HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J, H3FA; H3FL; H3FC; H3FB; H3FD; H3FI; H3FH; H3FK; H3FF; H3FJ
Immunogen Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Target Names	HIST1H3A

Image



Western Blot

Detected samples: HeLa whole cell lysate, Jurkat whole cell lysate, HepG2 whole cell lysate; Untreated (-) or treated (+) with 30mM sodium crotonylate for 4h


All lanes: HIST1H3A antibody at 1:100

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 16 kDa


Observed band size: 16 kDa



Immunocytochemistry analysis of CSB-PA010418OA09crHU diluted at 1:100 and staining in HeLa cells (treated with 30mM sodium crotonylate for 4h) performed on a Leica Bond™ system. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunoprecipitating HIST1H3A in HeLa whole cell lysate (treated with 30mM sodium crotonylate for 4h)
 Lane 1: Rabbit control IgG instead of CSB-PA010418OA09crHU in HeLa whole cell lysate (treated with 30mM sodium crotonylate for 4h). For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)
 Lane 2: CSB-PA010418OA09crHU (5µg) + HeLa whole cell lysate (treated with 30mM sodium crotonylate for 4h) (500µg)
 Lane 3: HeLa whole cell lysate (treated with 30mM sodium crotonylate for 4h) (20µg)



Chromatin Immunoprecipitation HeLa (10^6 , treated with 30mM sodium crotonylate for 4h) were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with 5µg anti-HIST1H3A (CSB-PA010418OA09crHU) or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers against the β -Globin promoter.

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.