



Phospho-RB1 (S780) Recombinant Monoclonal Antibody

Product Code	CSB-RA019386A780pHU
Abbreviation	Retinoblastoma-associated protein
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P06400
Immunogen	A synthesized peptide derived from Human Phospho-RB1 (S780)
Species Reactivity	Human
Tested Applications	ELISA, IHC, IF, IP; Recommended dilution: IHC:1:50-1:200, IF:1:20-1:200, IP:1:200-1:1000
Relevance	<p>Key regulator of entry into cell division that acts as a tumor suppressor. Promotes G0-G1 transition when phosphorylated by CDK3/cyclin-C. Acts as a transcription repressor of E2F1 target genes. The underphosphorylated, active form of RB1 interacts with E2F1 and represses its transcription activity, leading to cell cycle arrest. Directly involved in heterochromatin formation by maintaining overall chromatin structure and, in particular, that of constitutive heterochromatin by stabilizing histone methylation. Recruits and targets histone methyltransferases SUV39H1, KMT5B and KMT5C, leading to epigenetic transcriptional repression. Controls histone H4 'Lys-20' trimethylation. Inhibits the intrinsic kinase activity of TAF1. Mediates transcriptional repression by SMARCA4/BRG1 by recruiting a histone deacetylase (HDAC) complex to the c-FOS promoter. In resting neurons, transcription of the c-FOS promoter is inhibited by BRG1-dependent recruitment of a phospho-RB1-HDAC1 repressor complex. Upon calcium influx, RB1 is dephosphorylated by calcineurin, which leads to release of the repressor complex (By similarity). In case of viral infections, interactions with SV40 large T antigen, HPV E7 protein or adenovirus E1A protein induce the disassembly of RB1-E2F1 complex thereby disrupting RB1's activity.</p>
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Rabbit IgG in 10mM phosphate buffered saline , pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Purification Method	Affinity-chromatography
Isotype	Rabbit IgG
Clonality	Monoclonal
Alias	Exon 17 tumor GOS561 substitution mutation causes premature stop antibody; GOS563 exon 17 substitution mutation causes premature stop antibody; OSRC antibody; Osteosarcoma antibody; p105-Rb antibody; P105RB antibody; PP105 antibody; pp110 antibody; PPP1R130 antibody; pRb antibody; Prepro retinoblastoma associated protein antibody; Protein phosphatase 1 regulatory



subunit 130 antibody; Rb antibody; RB transcriptional corepressor 1 antibody; RB_HUMAN antibody; RB1 antibody; RB1 gene antibody; Retinoblastoma 1 antibody; Retinoblastoma susceptibility protein antibody; Retinoblastoma-associated protein antibody


Immunogen Species Homo sapiens (Human)

Research Area Cell Biology


Target Names RB1

Clone No. 2E9


Image



IHC image of CSB-RA019386A780pHUU diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4? overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of K562 cells with CSB-RA019386A780pHUU at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4?. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



Immunoprecipitating Phospho-RB1 in HeLa whole cell lysate
 Lane 1: Rabbit control IgG(1μg) instead of CSB-RA019386A780pHUU in HeLa whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)
 Lane 2: CSB-RA019386A780pHUU(3μg)+ HeLa whole cell lysate(1mg)
 Lane 3: HeLa whole cell lysate (20μg)

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.