




DBT Antibody

Product Code	CSB-PA006530LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P11182
Immunogen	Recombinant Human Lipoamide acyltransferase component of branched-chain alpha-keto acid dehydrogenase complex, mitochondrial protein (140-259AA)
Raised In	Rabbit
Species Reactivity	Human, Mouse
Tested Applications	ELISA, WB; Recommended dilution: WB:1:500-1:5000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Lipoamide acyltransferase component of branched-chain alpha-keto acid dehydrogenase complex, mitochondrial (EC 2.3.1.168) (52 kDa mitochondrial autoantigen of primary biliary cirrhosis) (Branched chain 2-oxo-acid dehydrogenase complex component E2) (BCOADC-E2) (Branched-chain alpha-keto acid dehydrogenase complex component E2) (BCKAD-E2) (BCKADE2) (Dihydrolipoamide acetyltransferase component of branched-chain alpha-keto acid dehydrogenase complex) (Dihydrolipoamide branched chain transacylase) (Dihydrolipoyllysine-residue (2-methylpropanoyl)transferase), DBT, BCATE2
Immunogen Species	Homo sapiens (Human)
Research Area	Tags & Cell Markers
Target Names	DBT

Image



Western Blot

Positive WB detected in: 293 whole cell lysate, Mouse kidney tissue, Mouse liver tissue

All lanes: DBT antibody at 4.3µg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 54 kDa

Observed band size: 54 kDa



Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.