


🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 📀 Website: www.cusabio.com 🌘

## SERPINB2 Antibody

Product Code	CSB-PA021070LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P05120
Immunogen	Recombinant Human Plasminogen activator inhibitor 2 protein (293-389AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	lgG
Clonality	Polyclonal
Alias	Plasminogen activator inhibitor 2 (PAI-2) (Monocyte Arg-serpin) (Placental plasminogen activator inhibitor) (Serpin B2) (Urokinase inhibitor), SERPINB2, PAI2 PLANH2
Immunogen Species	Homo sapiens (Human)
Research Area	Immunology
Target Names	SERPINB2
Image	Western Dist
	Western Blot




## Western Blot

Positive WB detected in: LO2 whole cell lysate, HepG2 whole cell lysate, Raji whole cell lysate All lanes: SERPINB2 antibody at 3.4µg/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 47 kDa Observed band size: 47 kDa

1



🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 📀 Website: www.cusabio.com 🍵



IHC image of CSB-PA021070LA01HU diluted at 1:200 and staining in paraffin-embedded human cervical cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.