






BROX Antibody

Product Code	CSB-PA730136LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q5VW32
Immunogen	Recombinant Human BRO1 domain-containing protein BROX protein (288-409AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500, IF:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	BRO1 domain-containing protein BROX (BRO1 domain- and CAAX motif-containing protein), BROX, BROFTI C1orf58
Immunogen Species	Homo sapiens (Human)
Research Area	Cell Biology
Target Names	BROX
_	





Western Blot

Positive WB detected in: K562 whole cell lysate

All lanes: BROX antibody at 3µg/ml


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 47, 43 kDa Observed band size: 47 kDa










IHC image of CSB-PA730136LA01HU diluted at 1:200 and staining in paraffin-embedded human pancreatic cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of Hela cells with CSB-PA730136LA01HU at 1:66, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).