




# MIB2 Antibody

|                            |   |
|----------------------------|---|
| <b>Product Code</b>        | CSB-PA839291LA01HU  |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.   |
| <b>Uniprot No.</b>         | Q96AX9  |
| <b>Immunogen</b>           | Recombinant Human E3 ubiquitin-protein ligase MIB2 protein (1-128AA)  |
| <b>Raised In</b>           | Rabbit  |
| <b>Species Reactivity</b>  | Human   |
| <b>Tested Applications</b> | ELISA, IHC; Recommended dilution: IHC:1:500-1:1000  |
| <b>Form</b>                | Liquid  |
| <b>Conjugate</b>           | Non-conjugated  |
| <b>Storage Buffer</b>      | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, pH 7.4  |
| <b>Purification Method</b> | >95%, Protein G purified  |
| <b>Isotype</b>             | IgG   |
| <b>Clonality</b>           | Polyclonal  |
| <b>Alias</b>               | E3 ubiquitin-protein ligase MIB2 (EC 2.3.2.27) (Mind bomb homolog 2) (Novel zinc finger protein) (Novelzin) (Putative NF-kappa-B-activating protein 002N) (RING-type E3 ubiquitin transferase MIB2) (Skeletrophin) (Zinc finger ZZ type with ankyrin repeat domain protein 1), MIB2, SKD ZZANK1 |
| <b>Immunogen Species</b>   | Homo sapiens (Human)  |
| <b>Research Area</b>       | Signal Transduction   |
| <b>Target Names</b>        | MIB2  |

## Image



IHC image of CSB-PA839291LA01HU diluted at 1:600 and staining in paraffin-embedded human skeletal muscle tissue performed on a Leica Bond<sup>TM</sup> system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.