





HACD3 Antibody

Product Code	CSB-PA878945LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9P035
Immunogen	Recombinant Human Very-long-chain (3R)-3-hydroxyacyl-CoA dehydratase 3 protein (1-145AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, IHC; Recommended dilution: IHC:1:200-1:500
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Very-long-chain (3R)-3-hydroxyacyl-CoA dehydratase 3 (EC 4.2.1.134) (3-hydroxyacyl-CoA dehydratase 3) (HACD3) (Butyrate-induced protein 1) (B-ind1) (hB-ind1) (Protein-tyrosine phosphatase-like A domain-containing protein 1), HACD3, BIND1 PTPLAD1
Immunogen Species	Homo sapiens (Human)
Research Area	Signal Transduction
Target Names	HACD3

Image



IHC image of CSB-PA878945LA01HU diluted at 1:200 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of CSB-PA878945LA01HU diluted at 1:200 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.